

AANBEVELINGEN TEST STRATEGIE

RAG subgroep testing – Update 14 December 2020

Opmerking: De huidige aanbevelingen zijn onderhevig aan veranderingen afhankelijk van nieuwe wetenschappelijke gegevens en/of de evolutie van de epidemie.

Voornaamste aanbevelingen:

Niveau 1 - Testen wordt als noodzakelijk beschouwd, en is dus sterk aanbevolen voor:

- Personen met suggestieve symptomen
- Hoog-risico contacten
- Laag-risico contacten tijdens een cluster onderzoek in scholen en bedrijven
- Terugkerende reizigers uit een rode zone
- Nieuwe bewoners van woon-zorgcentra (afhankelijk van de epidemiologische situatie)
- Niet-COVID patiënten bij opname in een risico afdeling van een ziekenhuis (afhankelijk van de epidemiologische situatie)
- Herhaald screenen van personeel van woon-zorgcentra en thuisverplegers (afhankelijk van de epidemiologische situatie).

Niveau 2 - Testen wordt als nuttig maar niet als noodzakelijk beschouwd, en is dus optioneel en afhankelijk van de epidemiologische situatie en andere voorwaarden waaraan moet worden voldaan:

- Bezoekers aan woon-zorgcentra
- Eenmalige screening van andere personen die potentieel veel anderen kunnen besmetten en waarbij effectieve preventieve maatregelen niet of moeilijk haalbaar zijn
- Herhaald screenen van andere bevolkingsgroepen die potentieel veel anderen kunnen besmetten en waarbij effectieve preventieve maatregelen niet of moeilijk haalbaar zijn.

Algemene aanbevelingen

- Een korte turn around time (doelstelling < 24u) blijft belangrijk, in alle testsituaties.
- De grootste prioriteit voor testen blijven de symptomatische personen en het onderzoek van clusters in collectiviteiten. Nu de test capaciteit het toelaat is het testen van hoog-risico contacten ook een prioriteit.
- Nieuwe technieken, zoals het afnemen van speekselstalen en snelle antigeentesten, creëren mogelijkheden voor nieuwe teststrategieën, met o.a. herhaaldelijk testen en eenmalige screening.
- Testen op zich volstaat niet om de epidemie te controleren. Alle test inspanningen hebben enkel zin als de preventieve maatregelen en de isolatie maatregel (en quarantaine voor contacten) correct nageleefd worden.
- Bepaalde test indicaties zijn enkel van toepassing in een context van hoge virus circulatie (= in functie van de epidemiologische situatie). Voor het bepalen van een drempelwaarde

hiervoor zal input van modelleringsonderzoek gebruikt worden. In afwachting worden de drempelwaarden voor de beleidsstrategie gehanteerd. In de controle fase (14-daagse cumulatieve incidentie < 100/100 000 en PR < 3%) is screening van asymptomatische personen, uitgezonderd hoge risico contacten en reizigers terugkerend uit een rode zone, minder zinvol.

- Door de toegenomen test capaciteit kunnen PCR testen en snelle Ag testen breder gebruikt worden, ook om socio-economische redenen, eerder dan volksgezondheid. De beslissing van wie voor een bepaalde test moet betalen moet door de politieke autoriteiten worden genomen, niet door de RAG.
- Belangrijk is wel dat (minstens) alle positieve resultaten van uitgevoerde testen via health/data en Sciensano aan de centra voor contactonderzoek worden gerapporteerd.
- Ook moet het uitvoeren van een test het wettelijk kader respecteren. Zo kan een snelle Ag test enkel uitgevoerd worden op voorschrift en onder de verantwoordelijkheid van een arts.
- Binnen het kader van dit advies werd geen exhaustieve lijst gemaakt van mogelijke toepassingen van gebruik van speekselstalen of snelle Ag testen voor screening, maar werden er wel criteria opgesteld voor de evaluatie van het nut van dergelijke screening. De RAG stelt voor om wekelijks de vragen voor adviezen rond screening te evalueren, aan de hand van een template document voor adviesaanvraag, dat opgesteld zal worden.

Gebruik van PCR-testen

PCR testen op een naso-faryngeale wisser of een gecombineerde keel-neuswisser blijven nog steeds de “gouden standaard” voor de diagnose van een infectie met SARS-CoV-2 (hoogste gevoeligheid). De capaciteit om PCR testen uit te voeren is ondertussen verhoogd wat mogelijkheden biedt voor een meer uitgebreid gebruik.

De PCR test blijft aangeraden daar waar er nog geen voldoende evidentie is dat het alternatief, de snelle antigeentest, voldoende betrouwbaarheid heeft. De test blijft aanbevolen voor de diagnose in symptomatische gevallen met een symptoomduur van meer dan 5 dagen, en bij patiënten op de spoed die onmiddellijk in het ziekenhuis moeten worden opgenomen. De test blijft ook aanbevolen voor het testen van hoog-risico contacten en reizigers terugkerend uit een rode zone, en voor herhaaldelijk testen op speekselstalen.

Net zoals in voorgaande RAG-adviezen wordt voor hoog-risicocontacten een eerste test aanbevolen zo snel mogelijk na de identificatie van het hoog-risico contact, met een maximum van 3 dagen na de laatste blootstelling. Deze test laat toe om snel positieve gevallen te detecteren en zo contactonderzoek rond deze personen te kunnen opstarten (backward contact tracing). Indien de contactpersoon pas laattijdig wordt geïdentificeerd (meer dan 72 uur na het laatste contact) is de eerste test niet meer nuttig en moet dus niet worden uitgevoerd. Daarnaast wordt, indien de quarantaine vroeger stop gezet wordt dan 10 dagen, een tweede test behouden ten vroegste 7 dagen na het laatste hoog-risicocontact, om infectie uit te sluiten vooraleer de quarantaine te beëindigen.

Reizigers die terugkeren uit een rode zone worden, net als de hoog-risico contacten, getest ten vroegste 7 dagen na hun terugkeer, om infectie uit te sluiten vooraleer de quarantaine te beëindigen, indien die vroeger stop gezet wordt dan 10 dagen.

Gebruik van multiplex PCR

De aanbeveling van de RAG in september blijft ongewijzigd. Multiplex PCR-testen, die tegelijkertijd meerdere respiratoire pathogenen detecteren, zijn nuttig tijdens de griep epidemie bij patiënten met ernstige respiratoire symptomen. De testen worden enkel aanbevolen in een hospitaalomgeving (patiënten op de spoedafdeling en gehospitaliseerde patiënten) en indien surveillancegegevens effectief een griep epidemie aantonen. Bij patiënten met zeer ernstige symptomen wordt aangeraden, indien mogelijk, een maxi panel te gebruiken (detectie van een breed scala aan pathogenen), bij patiënten met minder ernstige symptomen is een mini panel voldoende (enkel SARS-CoV2/ Influenza A/B, of enkel SARS-CoV2/ Influenza A/B en RSV).

Gebruik van snelle antigeen (Ag) testen

De snelle Ag testen van de tweede generatie zijn een nuttige aanvulling op de PCR-testen. Het blijft van cruciaal belang dat de selectie van de antigeentesten beperkt blijft tot de testen die voldoende gevalideerd zijn. Testen die, in patiënten met een symptoomduur van maximaal 5 dagen, een gevoeligheid van minder dan 80% of een specificiteit van minder dan 97% hebben (in vergelijking met een RT-PCR) mogen niet gebruikt worden. De gewenste gevoeligheid is >90%, en in personen met een hoge virale lading ($\geq 10^5$ RNA kopieën/mL of Ct waarde <25), die sterk besmettelijk zijn, moet het >95% zijn. Deze drempelwaarden moeten bevestigd zijn door ten minste drie onafhankelijke evaluaties.

Er zijn reeds testen beschikbaar die aan de voorwaarden hierboven voldoen en die worden aanbevolen bij symptomatische personen met klachten van een luchtweginfectie sedert ≤ 5 dagen in een huisartsenpraktijk, triagecentrum of spoeddienst - zie [RAG advies November](#).

Momenteel zijn er nog weinig wetenschappelijke gegevens over de gevoeligheid van de snelle Ag testen bij asymptomatische personen, maar de testen laten wel toe om hoog besmettelijke personen ($\geq 10^5$ RNA kopieën/mL of Ct waarde <25) te identificeren. Ag testen worden daarom ook aanbevolen voor een onderzoek van een cluster, waar de verwachte prevalentie hoger is en een snelle detectie bijdraagt tot het onmiddellijk nemen van de gepaste maatregelen om de cluster in te dijken. Momenteel werden reeds protocollen uitgewerkt voor het onderzoek van clusters in middelbare scholen en bedrijven. Voor een onderzoek van een cluster in woonzorgcentrum (WZC) worden de resultaten van de lopende studie in deze setting afgewacht.

Snelle Ag testen kunnen ook gebruikt worden voor het screenen van (asymptomatische) personen die eenmalig potentieel veel anderen kunnen besmetten (relatief hoge prevalentie, en nauw contact met een hoog aantal personen of verdere verspreiding mogelijk binnen een collectiviteit) of in contact komen met personen met een risico op een ernstige infectie; en waarbij effectieve preventieve maatregelen niet of moeilijk haalbaar zijn. Zo kan een WZC overwegen om screening aan te bieden aan bezoekers, om te voorkomen dat een positief en sterk besmettelijk persoon de kwetsbare bewoners besmet. Een negatief resultaat sluit echter niet uit dat de bezoeker alsnog infectieus is, en de geldende voorzorgsmaatregelen moeten daarom nog steeds in acht genomen worden. Ook geldt het resultaat van een snelle Ag test enkel voor de dag zelf. Voor personen die meermaals per week op bezoek gaan in het WZC kan overwogen worden om over te gaan tot een herhaalde screening (zie verder).

Het gebruik van snelle Ag testen voor kinderen jonger dan 12 jaar in het kader van een eenmalige screening wordt niet aanbevolen. Het afnemen van een neus/keel of naso-

faryngeaal staal moet in deze groep voorbehouden worden voor het vaststellen van een besmetting of voor hoog-risicocontacten, en enkel in kinderen ≥ 6 jaar.

Enmalige screening wordt momenteel niet sterk aangeraden, maar kan overwogen worden indien het vanuit een volksgezondheidstandpunt verantwoord is. Screening zou nooit als hoofddoel een versoepeling van de maatregelen mogen hebben. Indien de maatregelen versoepeld worden om andere, socio-economische redenen, kan een systematisch screenen wel een middel zijn om de negatieve gevolgen van deze versoepelingen tot een minimum te beperken.

Gebruik van speekselstalen

- Speekselstalen kunnen enkel onderzocht worden op de aanwezigheid van SARS-CoV-2 aan de hand van een PCR. Er lopen studies over het gebruik van de snelle antigeentesten op speeksel, maar dergelijke combinatie is tot op heden onvoldoende gevalideerd.
- Er is nog steeds onvoldoende wetenschappelijke evidentie om het gebruik van speekselstalen als diagnostische tool bij een symptomatische persoon aan te bevelen of om ze te gebruiken bij kinderen. Enkel indien een afname van aan naso-faryngeale wisser of een gecombineerde keel-neuswisser niet mogelijk of moeilijk is kan een speekselstaal overwogen worden.
- Een protocol werd opgesteld voor de afname en analyse van speekselstalen¹.
- De huidige PCR capaciteit laat het gebruik van speekselstalen in een setting van herhaald screenen bij asymptomatische personen (≥ 12 jaar) toe.
- Gezien de voorgestelde voorwaarden voor het starten ervan vervuld zijn (protocol en voldoende PCR capaciteit), kan herhaald (minstens wekelijks) screenen via speekselstalen nu aanbevolen worden bij personeel van WZC en bij thuisverplegers van personen kwetsbaar voor een ernstige COVID-19 infectie.
- Herhaald screenen van andere populaties wordt momenteel niet aangeraden, maar kan wel overwogen worden indien het vanuit een volksgezondheidstandpunt verantwoord is, namelijk bij personen die potentieel veel anderen kunnen besmetten (relatief hoge prevalentie, en nauw contact met een hoog aantal personen of verdere verspreiding mogelijk binnen een collectiviteit) of in contact komen met personen met een risico op een ernstige infectie; en waarbij effectieve preventieve maatregelen niet of moeilijk haalbaar zijn. Herhaald screenen zou nooit als hoofddoel een versoepeling van de maatregelen mogen hebben. Indien de maatregelen versoepeld worden om andere, socio-economische redenen, kan herhaaldelijk testen wel een middel zijn om de negatieve gevolgen van deze versoepelingen tot een minimum te beperken.
- Omdat de precieze accuraatheid nog niet helemaal gekend is wordt het gebruik van speekseltesten momenteel niet aanbevolen in andere omstandigheden, zoals voor het testen van nauwe contacten of terugkerende reizigers.

¹ Beschikbaar op: https://covid-19.sciensano.be/sites/default/files/Covid19/20201130_Advice%20RAG_Saliva%20sampling_NL.pdf

De volgende personen hebben deelgenomen aan dit advies:

Emmanuel André (KULeuven); Emmanuel Bottieau (ITG); Bénédicte Delaere (CHU-UCL Namur); Olivier Denis (CHU-UCL Namur); Frédéric Frippiat (AViQ); Herman Goossens (UAntwerpen); Marie Pierre Hayette (CHU-Liège); Niel Hens (UHasselt); Yves Lafort (Sciensano); Barbara Legiest (ZG); Tinne Lernout (Sciensano); Pieter Libin (UHasselt); Romain Mahieu (COCOM); Elizaveta Padalko (UZGent); Petra Schelstraete (UGent); Olivier Vandenberg (LHUB-ULB); Ann Van den Bruel (KuLeuven); Dimitri Van der Linden (UCLouvain); Pieter Vermeersch (UZ-Leuven).

Samenvatting test aanbevelingen

Indicatie	Aanbevolen staal	Aanbevolen test	Type aanbeveling
Symptomatische persoon			
Symptomen ≤ 5 dagen	Naso-faryngeale of keel-neuswisser	Ag RDT	In triage centra, huisartspraktijk ² en spoedgevallendienst
Symptomen > 5 dagen of dringende hospitalisatie	Naso-faryngeale of keel-neuswisser	PCR	Algemeen aanbevolen
Ernstig ziek tijdens griepiepidemie	Naso-faryngeale of keel-neuswisser	Multiplex PCR Mini-panel	Enkel in een ziekenhuis omgeving
Zeer ernstig ziek tijdens griepiepidemie	Naso-faryngeale of keel-neuswisser	Multiplex PCR Maxi-panel	Enkel in een ziekenhuis omgeving
Asymptomatische hoog-risico contacten			
Hoog-risico contact, direct na identificatie	Naso-faryngeale of keel-neuswisser	PCR	Indien laatste contact ≤ 3 dagen
Hoog-risico contact, 7 dagen na laatste contact	Naso-faryngeale of keel-neuswisser	PCR	Indien quarantaine vroegtijdig (<10 dagen) gestopt wordt
Asymptomatische reizigers			
Terugkerende reiziger uit rode zone (dag 7)	Naso-faryngeale of keel-neuswisser	PCR	Indien quarantaine vroegtijdig (<10 dagen) gestopt wordt
Cluster onderzoek			
Lage risico contacten in school cluster	Naso-faryngeale of keel-neuswisser	Ag RDT	Enkel in middelbare scholen
Lage risico contacten in bedrijf cluster	Naso-faryngeale of keel-neuswisser	Ag RDT	Alle clusters in een bedrijfsomgeving
Herhaald screenen			
Herhaald testen personeel WZC en thuisverplegers	Speeksel	PCR	Aanbevolen in functie van epidemiologische situatie
Herhaald testen andere populaties	Speeksel	PCR	Optioneel en enkel indien aan bepaalde voorwaarden voldaan
Enmalige screening			
Ziekenhuisopname niet-COVID patiënt	Naso-faryngeale of keel-neuswisser	PCR	Volgens de bestaande richtlijnen voor ziekenhuizen ³
Nieuwe bewoners WZC	Naso-faryngeale of keel-neuswisser	PCR	Aanbevolen in functie van epidemiologische situatie
Bezoekers WZC	Naso-faryngeale of keel-neuswisser	Ag RDT	Optioneel
Andere situaties	Naso-faryngeale of keel-neuswisser	Ag RDT	Optioneel en enkel indien aan bepaalde voorwaarden voldaan

² Van zodra de lopende pilootstudies beëindigd zijn en indien deze een positieve uitkomst hadden

³ Beschikbaar op: https://covid-19.sciensano.be/sites/default/files/Covid19/Brief_Teststrategie%20voor%20algemene%20en%20psychiatrische%20ziekenhuizen%20en%20voor%20de%20revalidatieziekenhuizen.pdf

CONTEXT

The PCR test capacity has substantially improved over the past month, better performing rapid antigen tests have become available and a protocol for sampling and analyzing saliva specimens has been developed. This opened perspectives for new testing strategies.

The current test strategy update focusses on the following topics:

1. Testing of asymptomatic high-risk contacts	7
2. Repetitive screening.....	10
3. One-time screening.....	15
4. Use of multiplex PCR assays	18

1. TESTING OF ASYMPTOMATIC HIGH-RISK CONTACTS

1.1. Definition of a high-risk contact

The current definition of a contact person is any person who has had contact with a confirmed COVID-19 case within a period of 2 days before the onset of symptoms (symptomatic cases) or before the sample was taken (asymptomatic cases) until the end of the infectivity period of the case - generally 7 days after the start of the symptoms, or longer if symptoms persist, (symptomatic cases) or 7 days after the sample was taken (asymptomatic cases).

A contact person is considered a high-risk contact if any of the following conditions are present⁴:

- a cumulative contact of at least 15 minutes within a distance of <1.5 m ("face to face") without proper use of a mouth mask by either person;
- direct physical contact with a COVID-19 patient;
- direct physical contact with excretions or body fluids of a COVID-19 patient;
- been identified by the "Coronalert" application as a close contact;
- traveled with a COVID-19 patient for more than 15 minutes.

1.2. Past and current recommendations

Recommendations for testing (asymptomatic) high-risk contacts (HRCs) have changed over time in response to the evolution of the epidemic and the available testing capacity. Initially, only HRCs with a high risk of coming in contact with vulnerable populations were tested (with a PCR) at the end of the 14-days quarantine period (between day 11 and 13), to confirm that they are not infected.

In June (12/6), when the incidence had decreased and the contact tracing system had become operational, the recommendation became to test all HRCs, except children under 6, immediately after identification to detect positive cases. HRCs testing negative went into a 14 days quarantine period. The rule to test HRCs in contact with vulnerable populations at the end of the quarantine period was maintained.

At the end of June (23/6), the protocol was revised. If the test was done within 7 days of the last contact, a second test could be performed the earliest 5 days after the first test and 9 days

⁴ More detailed definition available at: https://covid-19.sciensano.be/sites/default/files/Covid19/COVID-19_procedure_contact_NL.pdf

after the last contact. If both tests were negative, the quarantine could be stopped at day 10. HRCs in contact with vulnerable populations could also be tested in the first week after exposure, but had to remain in quarantine if testing negative. Only if testing negative on day 11 to 13, the quarantine could be stopped on day 14.

At the beginning of October (1/10), when incidence had increased again and the 14-day quarantine period resulted in some staff shortages, the strategy changed to no longer test HRCs immediately after identification and only test them the earliest 5 days after the last contact. If testing negative, the quarantine could be stopped on day 7.

At the end of October (21/10), testing of asymptomatic HRCs was put on a hold because of the high pressure on the PCR testing capacity.

Since November 23, testing of asymptomatic HRCs has reinitiated. The current recommendation is the same as the October 1st recommendation:

- All asymptomatic close contacts (except children <6 years) should be tested at the earliest 7 days after the last high-risk contact.
 - If the result is positive, the home isolation is extended to 7 days after the test (or start of symptoms). The contact becomes an index case and the detection of his or her closest contacts is initiated.
 - If the result is negative, the quarantine can be lifted as soon as the test result is obtained (minimum 7 days after the last high-risk contact). As the incubation period can be up to 14 days, the general measures must be followed for a further 7 days.

1.3. International recommendations and strategies in other countries

ECDC

ECDC updated its recommendations with regards to contact tracing on November 18⁵.

It recommends that high-risk contacts without symptoms are tested as soon as possible after being traced, to enable early identification of any asymptomatic or pre-symptomatic secondary cases among contacts and to start further contact tracing. This also applies to low-risk contacts in settings with vulnerable populations or in which transmission is likely, such as health and social care settings, prisons, certain occupational settings and social events such as choirs or weddings.

Both RT-PCR and Ag RDTs can be considered. Ag RDTs have the advantage of speed which allows for further contact tracing to commence as soon as possible. If more than seven days have passed since the exposure, it is recommended that negative Ag RDTs are confirmed by RT-PCR.

Even if the test is negative, high-risk exposure contacts still need to quarantine, and low-risk exposure contacts still need to observe prevention measures for the remainder of the 14 days after exposure.

Testing at the end of quarantine can also be done, in addition to the test taken upon tracing. A negative RT-PCR test at day 10 can be used to discontinue quarantine earlier than the recommended 14 days.

Testing capacity may be limited and it is important that testing contact persons for the purpose of ending quarantine early should not adversely impact test accessibility and test turnaround time for symptomatic people.

⁵ ECDC - TECHNICAL REPORT - Contact tracing: public health management of persons, including healthcare workers, who have had contact with COVID-19 cases in the European Union – third update - 18 November 2020

In the Technical Report for the use of rapid antigen tests (November 19), ECDC recommends to only test contacts with a Ag RDTs if the test positivity is expected to be $\geq 10\%$ ⁶.

WHO

WHO has not updated its guidance on contact tracing since August 19. The latest guidance only stipulated that for contacts who do not develop symptoms, WHO no longer considers laboratory testing a requirement for leaving quarantine after 14 days.

France

In France, asymptomatic contacts are advised to immediately get tested if they live under the same roof as the index case⁷. If they do not live under the same roof, the advice is to get tested 7 days after the last contact. The rationale is that testing earlier has a too great chance of giving a negative result. Contacts living under the same roof testing negative in the first test have to test again 7 days after the index case is cured. If the test is negative, the quarantine can be stopped.

Other neighboring countries

The Netherlands and England do not recommend to test contacts. No guidelines were found for Germany.

1.4. Discussion

- A first test early after identification enables early detection of any asymptomatic or pre-symptomatic secondary cases among contacts and starting further contact tracing.
- All HRCs ≥ 6 years old should be tested, including those not living under the same roof, because these might be the index case.
- Systematically testing of low risk contacts in settings with vulnerable populations is not indicated. The current guideline is that the health inspector evaluates the situation when a case is detected in a collectivity with a vulnerable population, such as a nursing home, and can decide, if the risk of an outbreak is high, to test low-risk contacts. This guideline remains adequate.
- It will be difficult to return to a longer quarantine period, because this would face too much resistance of the public, the industry and politicians, and the current recommended period of minimum 7 days should be maintained. A second test on day 7 after the last contact remains therefore indicated. However, if the high-risk contact is only identified several days after the last contact, the period between the first and second test may become very short. There should be at least 4 days between the two tests. In this event, if the contact is identified more than three days after the last contact with the index case, it is indicated to skip the first test and only test on day 7.
- A separate quarantine and testing procedure for contacts living under the same roof and contacts not living under the same roof, as in France, makes it too complicated and is not necessary. The current procedure already takes into account situations where it is not possible to isolate household contacts from each other (such as between parents and small children) and considers in those situations the last day of the 7 day isolation period as the last contact.

⁶ ECDC - TECHNICAL REPORT - Options for the use of rapid antigen tests for COVID-19 in the EU/EEA and the UK - 19 November 2020. Available at : <https://www.ecdc.europa.eu/en/publications-data/options-use-rapid-antigen-tests-covid-19-eueea-and-uk>

⁷ J'ai été en contact à risque avec une personne malade du COVID-19 [Fiche patients] - Publié le 3 Novembre 2020 - Mis à jour le 06 novembre 2020. Available at : <https://www.santepubliquefrance.fr/maladies-et-traumatismes/maladies-et-infections-respiratoires/infection-a-coronavirus/documents/depliant-flyer/j-ai-ete-en-contact-a-risque-avec-une-personne-malade-du-covid-19-fiche-patients>

- In both the first and the second test the more sensitive PCR test is indicated, because the contact might still be in an early stage of infection before viral load increases. This also applies to the second test, because the current quarantine period is much shorter than the incubation period.
- In periods that there is pressure on the PCR capacity and long delays in getting the results (>24h hours), an Ag RDT might be considered as an alternative.
- Testing high-risk contacts is only effective if the contact tracing system is effective, which is currently still insufficiently the case. There is therefore a need to further improve the system.

1.5. Recommendations

- To test all HRCs, 6 years old or older, as soon as possible after their identification.
- To re-test HRCs, who tested negative in the first test, on day 7 after their last contact with the index patient
- To skip the first test if the HRC is identified more than 3 days after the last contact. In that event the HRC is only tested on day 7.
- To use a PCR test for both the first and second test episode, unless there is a delay of more than 24 hours in getting the result. In that event an Ag RDT can be used.
- Low-risk contacts are only tested in the context of a cluster investigation, or when the health inspector judges that there is a risk of an outbreak in a collectivity with a vulnerable population (see separate procedures).

2. REPETITIVE SCREENING⁸

2.1. Current recommendation

The current recommendation (November update) with regards to repetitive screening (using saliva samples) is:

- There is an urgent need to establish a protocol for the use of repetitive saliva samples in combination with PCR in various sectors (health care, education, high-risk companies, essential services).
- If there is an increased demand for saliva testing in a setting of repetitive screening, the pressure on PCR testing capacity will further increase. Therefore, before starting repeated screening using saliva samples, the capacity must first be increased.
- If the two conditions mentioned above (sufficient PCR capacity and protocol available) are met, the use of saliva samples for repeated screening (at least weekly) in asymptomatic adults, such as staff of nursing homes, can be considered.

2.2. Literature

Several modelling studies have assessed the effectiveness of repetitive screening in controlling SARS-CoV-2 spread in specific populations. All these models showed that frequent testing with a less sensitive test (rapid antigen test) or a less sensitive sample (saliva) is more effective than one-time testing with the more sensitive RT-PCR on a naso-pharyngeal sample

⁸ For the purpose of this advice we define 'repetitive screening' as testing asymptomatic people at frequent regular intervals to rapidly detect an infection by SARS-CoV-2.

(1–3). The recommended periodicity varies across models and is dependent on various factors, such as the sensitivity and specificity of the test, the prevalence, the reproductive number and the compliance to measures taken for positive cases. Most studies recommend a periodicity of at least 2-3 times a week (4–7), but other state that relatively infrequent testing, such as every one or two weeks, is already sufficient to keep controlled outbreaks small (8). One study assessed the effect of regular universal testing and concluded that this strategy would require unrealistic high testing frequencies to reopen society while maintaining control of virus transmission (9).

Atkeson et al. assessed the economic benefits of repeated testing (with a rapid antigen test) and concluded that the fiscal, macroeconomic, and health benefits of rapid SARS-CoV-2 screening testing programs far exceed their costs (10). A weekly testing in a regime with high compliance comes close to suppressing the virus, and moving to a four-day cadence is highly effective. They point out however, that the screening testing program must have high specificity to be credible and to evoke high adherence. If specificity is not close to 100%, the positive predictive value is low in low-prevalence settings, putting many people unnecessary in isolation. They propose therefore confirmation of positive results with an RT-PCR test. The problem of low positive predictive value in low prevalence settings, and therefore a need to confirm positive results, is also addressed in another study (11).

Studies evaluating the effect of repetitive screening in a real-life situation are rare. One study evaluated a longitudinal screening program for critical on-site employees within a research institute, and concluded that it was accepted by employees and can be used to maintain the health of the workforce, potentially keeping positivity rates below community levels (12). Another study screened asymptomatic HCWs in a large hospital over a 3-week period and concluded that such an approach is critical for protecting patients and hospital staff (13). Both these studies used RT-PCR tests.

2.3. International recommendations

Very few countries have issued guidelines on repetitive screening. **ECDC** mentions in their '*Population-wide testing of SARS-CoV-2: country experiences and potential approaches in the EU/EEA and the United Kingdom*' report (19 August 2020) that most EU/EEA countries and the UK regularly test individuals in high-risk settings such as at healthcare facilities⁹. Of the eight countries that responded to the ECDC enquiry that are not currently planning population-wide testing of individuals without symptoms, at least five regularly test individuals without any symptoms in high-risk settings e.g. healthcare workers, individuals working in long-term care facilities, and people in various other settings, such as patients admitted to hospitals, individuals in specific occupational settings, prisons, etc. Testing in these settings was sometimes occurring in response to a cluster of cases reported in that group or setting, or carried out to protect vulnerable populations (i.e. patient groups, health workers, patients or long-term care facility residents).

WHO

In its interim guidance on Prevention, identification and management of health worker infection in the context of COVID-19 (30 October 2020), the World Health Organization lists factors that should be taken into account when deciding on the need for routine testing among health care workers¹⁰:

⁹ Available at: <https://www.ecdc.europa.eu/sites/default/files/documents/covid-19-population-wide-testing-country-experiences.pdf>

¹⁰ Available at: [file:///C:/Users/YvLa1869/Downloads/WHO-2019-nCoV-HW_infection-2020.1-eng%20\(2\).pdf](file:///C:/Users/YvLa1869/Downloads/WHO-2019-nCoV-HW_infection-2020.1-eng%20(2).pdf)

- The intensity of transmission in the setting of the health facility(ies), for example in the presence of community transmission or intense outbreaks of COVID-19.
- The capacities of the facility and laboratories to conduct the testing including financial and human resources available, as well as availability of testing materials and laboratory capacity.
- The volume of patients identified as positive for SARS-CoV-2, admitted to the facility or being assessed by health workers.
- The positivity rate among staff.
- The number of staff who are ill but not diagnosed with COVID-19 and in quarantine as contacts for COVID-19, leading to inability to provide adequate and safe staffing levels.

For health workers staffing or working with long-term care facilities, they recommend routine testing irrespective of the COVID-19 transmission scenario.

CDC

The US Centers for Disease Control recommends that staff of nursing homes be serially tested based on the county percent test positivity rate. Approaches may include initial testing of all workers before entering a workplace, periodic testing of workers at regular intervals, and/or targeted testing of new workers or those returning from a prolonged absence.

2.4. Discussion

- The main objective of repetitive screening should be a public health one, namely to early detect positive infections in people whose infection can have serious public health consequences (potentially infect many other people or infect vulnerable people).
- The objective of repetitive screening is not to allow relaxation of preventive measures, or to avoid isolation or quarantine. Nevertheless, if epidemic control measures are alleviated for socio-economic reasons, repetitive screening can be a mean to mitigate the negative impact of these relaxations.
- Lower sensitivity of the test used can be compensated by a sufficiently high frequency.
- The use of saliva samples has the advantage of ease of sample taking, but requires the more expensive RT-PCR.
- The use of rapid antigen tests has the advantage of the lower cost and rapidity in getting results, but requires nasopharyngeal, or at least nasal/oral, specimens.

Criteria to consider when deciding on repetitive screening in a specific population (in order of importance):

- Having contact with people vulnerable to severe COVID-19 disease
Rapid detection of positive COVID-19 cases is especially important in people who are continuously in contact with persons vulnerable to a severe form of the disease. These are in the first place elderly people. Staff in nursing homes and home nurses attending to elderly at their home are therefore populations eligible for repetitive screening.
- Not being able to take the appropriate protective measures
Repetitive screening is less meaningful in populations that are perfectly able to correctly protect themselves and the people they come in contact with.
- Potential to rapidly transmit the infection to a large number of people
Repetitive screening is only meaningful in populations that, if infected, have a potential to rapidly transmit the infection to a large number of people. These can be people who have

a large number of serial close contacts, are come in close contact with large groups of people, or live in social clusters, and who have a substantial risk to be infected.

- Feasibility
Before initiating repetitive screening in a specific population the feasibility needs to be assessed. This is especially important if the size of the population is large.

Criteria to consider in the choice of the specimen to collect and the test to use:

- Ease of sampling
Because frequent sampling is inherent to repetitive screening, it is recommended to use as much as possible specimens that are little invasive or painful, and easy to collect. Nasopharyngeal samples are therefore not recommended. Nasal/oral swabs are acceptable, but saliva is by far the least invasive specimen.
- Cost
The rapid antigen test is much less expensive than an RT-PCR. Nevertheless, the current PCR capacity allows to perform a large number of tests at relatively little additional cost.
- Rapidity of result
In settings where a very rapid result is required, a rapid antigen test is indicated. However, in a context of sufficient RT-PCR capacity, the turn-around time to get the PCR result should be sufficiently short in most situations.

Testing frequency

The ideal testing frequency depends on a number of factors, such as the sensitivity of the testing protocol used, the positivity rate in the tested population, the incidence and the reproductive number, and the measures taken for positive cases (and level of adherence to those measures). A European consortium, with UHasselt as one of the members, is currently developing a mathematical model that will provide more exact recommendations on the ideal testing frequency in different situations and prevalence levels. Pending the results, we maintain the recommendation to test at least weekly.

Result interpretation

A sufficiently high frequency of testing compensates for the lower sensitivity of a saliva specimen or an Ag RDT, and a person with a negative test result is therefore considered as not having COVID-19 and/or not being infectious.

The use of RT-PCR guarantees a sufficiently high specificity. If an Ag RDT is used with a lesser specificity, there is a risk of a low positive predictive value if the positivity rate in the tested population is very low. However, testing in populations with a very low prevalence is anyway not indicated and even then have false positive detections few negative consequences.

2.5. Recommendations

- We recommend repetitive screening in populations that are in frequent contact with people vulnerable to severe COVID-19 disease (staff at nursing homes, home nurses attending to elderly clients).
- Repetitive screening in other populations is only justifiable if the following conditions are fulfilled:
 - Risk of rapid spread of the infection to a large number of people; AND
 - No possibility to fully apply effective protective measures; AND

- The number of people to test is feasible.
- Frequency of testing will be defined by the modelling exercise currently in development. Meanwhile, we recommend a frequency of at least weekly.
- We recommend the use of an RT-PCR test on saliva specimens. The use of Ag RDTs on nasal/oral swabs can be a valid alternative if the delay in getting the result is >24 hours because of reduced RT-PCR capacity.
- People testing negative are considered as not having COVID-19 and/or not being infectious. Basic preventive measures need nevertheless to be maintained. People testing positive are considered as confirmed cases and the procedures with regards isolation and contact tracing is initiated.

References

1. Mina MJ, Parker R, Larremore DB. Rethinking Covid-19 Test Sensitivity — A Strategy for Containment. *N Engl J Med* [Internet]. 2020 Sep 30 [cited 2020 Oct 15]; Available from: <https://www.nejm.org/doi/10.1056/NEJMp2025631>
2. Larremore DB, Wilder B, Lester E, Shehata S, Burke JM, Hay JA, et al. Test sensitivity is secondary to frequency and turnaround time for COVID-19 surveillance. *medRxiv*. 2020 Sep 8;2020.06.22.20136309.
3. Plebani M, Aita A, Cattelan AM, Bonfante F, Padoan A, Giaquinto C, et al. Frequent testing regimen based on salivary samples for an effective COVID-19 containment strategy. *medRxiv*. 2020 Oct 14;2020.10.13.20210013.
4. Hellewell J, Russell TW, Team TSI and FS, Consortium TCC-19, Group CC-19 working, Beale R, et al. Estimating the effectiveness of routine asymptomatic PCR testing at different frequencies for the detection of SARS-CoV-2 infections. *medRxiv*. 2020 Nov 24;2020.11.24.20229948.
5. Holmdahl I, Kahn R, Hay J, Buckee CO, Mina M. Frequent testing and immunity-based staffing will help mitigate outbreaks in nursing home settings. *medRxiv*. 2020 Nov 23;2020.11.04.20224758.
6. Chin ET, Lo NC, Huynh BQ, Murrill M, Basu S. Frequency of routine testing for SARS-CoV-2 to reduce transmission among workers. *MedRxiv Prepr Serv Health Sci*. 2020 May 6;
7. Chang JT, Crawford FW, Kaplan EH. Repeat SARS-CoV-2 Testing Models for Residential College Populations. *medRxiv*. 2020 Jul 16;2020.07.09.20149351.
8. Nash B, Badea A, Reddy A, Bosch M, Salcedo N, Gomez AR, et al. The impact of high frequency rapid viral antigen screening on COVID-19 spread and outcomes: a validation and modeling study. *medRxiv*. 2020 Nov 4;2020.09.01.20184713.
9. Bootsma MCJ, Kretzschmar ME, Rozhnova G, Heesterbeek J a. P, Kluytmans J, Bonten MJM. Regular universal screening for SARS-CoV-2 infection may not allow reopening of society after controlling a pandemic wave. *medRxiv*. 2020 Nov 18;2020.11.18.20233122.
10. Atkeson A, Droste M, Mina MJ, Stock JH. Economic Benefits of COVID-19 Screening Tests. *medRxiv*. 2020 Nov 1;2020.10.22.20217984.
11. Sudlow C, Diggle P, Warlow O, Seymour D, Gordon B, Walker R, et al. Testing for coronavirus (SARS-CoV-2) infection in populations with low infection prevalence: the largely ignored problem of false positives and the value of repeat testing. *medRxiv*. 2020 Aug 22;2020.08.19.20178137.

12. Goetz LH, DeLaughter TL, Kennedy KL, Schork NJ, McDaniel TK, Trent JM, et al. Asymptomatic Employee Screening for SARS-CoV-2: Implementation of and Reactions to an Employer-Based Testing Program. medRxiv. 2020 Nov 10;2020.11.06.20227314.

13. Rivett L, Sridhar S, Sparkes D, Routledge M, Jones NK, Forrest S, et al. Screening of healthcare workers for SARS-CoV-2 highlights the role of asymptomatic carriage in COVID-19 transmission. medRxiv. 2020 May 15;2020.05.09.20082909.

3. ONE-TIME SCREENING¹¹

3.1. Current recommendation

Testing of asymptomatic people is currently (RAG advice of November 3) advised in the context of contact tracing, returning travelers, cluster investigations and repetitive screening of nursing home staff.

One-time screening is only advised for non-COVID patients requiring hospitalization, under certain conditions, and new residents of nursing homes. The test to use is an RT-PCR. If the test is negative, it can be repeated according to clinical need.

3.2. Literature

Few studies assessing the effect of routine one-time screening in specific situations were identified.

A Cochrane review of universal screening for SARS-CoV-2 infection, done in September, concluded that the evidence base for the effectiveness of screening is limited. Low-certainty evidence suggests that screening at travel hubs may slightly slow the importation of infected cases. The authors highlight the uncertainty and variation in accuracy of screening strategies. A high proportion of infected individuals may be missed and go on to infect others, and some healthy individuals may be falsely identified as positive, requiring confirmatory testing and potentially leading to the unnecessary isolation of these individuals. Further studies need to evaluate the utility of rapid laboratory tests, combined screening, and repeated screening. More research is also needed on reference standards with greater accuracy than RT-PCR (1).

An update in October, concluded that one-time screening in apparently healthy people is likely to miss people who are infected. The authors are unsure whether combined screenings, repeated symptom assessment, or rapid laboratory tests are useful (2).

3.3. International recommendations

WHO

In their Interim guidance for antigen-detection in the diagnosis of SARS-CoV-2 infection using rapid immunoassays of 11 September 2020, WHO does not recommend the use of rapid antigen tests in individuals without symptoms unless the person is a contact of a confirmed case, and neither for airport or border screening at points of entry¹².

¹¹ For the purpose of this advice we define 'one-time screening' as testing asymptomatic people prior to a potential risk of spreading SARS-CoV-2 to others. Screening can occur in the same individual on different occasions, but unlike repetitive screening it is not done on pre-defined regular intervals.

¹² Available at: file:///C:/Users/YvLa1869/Downloads/WHO-2019-nCoV-Antigen_Detection-2020.1-eng%20(1).pdf

ECDC

In the guidelines for the use of rapid antigen tests (19 November 2020) ECDC states: 'In a high prevalence situation, in the context of circuit breaker strategies to detect individuals with high transmission potential in the community and to lower the pressure on health-care settings and laboratories, rapid antigen tests' use can be considered for a targeted population-wide testing approach, e.g. in a local community. In such situation, the risk of not detecting all cases or risk of false negative results is balanced out by the timeliness of results and the possibility of serial testing of individuals.' and 'Rapid antigen tests are not suited for screening incoming travelers to prevent virus (re-) introduction in regions/countries that have achieved zero or very low levels of transmission. In these situations, i.e. in a low prevalence population, only RT-PCR should be used to reduce the risk of false negative results. Air travelers belong mostly to a non-symptomatic sub-population, with variable but lower probability of COVID-19 compared to the general population (estimated prevalence of COVID-19 at <1%)¹³.

3.4. Discussion

- The main objective of one-time screening should be a public health one, namely to prevent an infected person to infect others before an at-risk exposure.
- The objective of one-time screening is not to allow relaxation of preventive measures, or to avoid isolation or quarantine. Nevertheless, if epidemic control measures are alleviated for socio-economic reasons, preventive screening can be a mean to mitigate the negative impact of these relaxations.
- Screening is less relevant in children <12 years, because the lesser role they play in spreading the SARS-CoV-2 virus..

Criteria to consider when deciding on routine one-time screening in a specific situation (in order of importance):

- People entering in contact with people vulnerable to severe COVID-19 disease
These are in the first place elderly people. Routine testing of visitors to nursing homes is therefore a meaningful strategy.
- Not being able to take the appropriate protective measures
Screening is less meaningful in situations where it is perfectly possible to correctly prevent transmission.
- Important contribution to the chain of transmission
Screening is most meaningful in situations where an early detection of positive cases can prevent further spread of the virus. It is therefore most useful in populations with a high risk of being infected who enter in contact with many other people, or enter in contact with a person in contact with many other people.
- Feasibility
Before initiating routine screening in a specific setting the feasibility needs to be assessed. Factors to consider with regards to feasibility include the amount of screening tests to perform and the feasibility to perform a rapid antigen test on site.

¹³ Available at : <https://www.ecdc.europa.eu/sites/default/files/documents/Options-use-of-rapid-antigen-tests-for-COVID-19.pdf>

Criteria to consider in the choice of the specimen to collect and the test to use

- Rapidity of result
In situations where a very rapid result is required, a rapid antigen test is indicated. This is especially the case when screening is done at the moment of entry.
- Cost
The low cost of the rapid antigen test is an important advantage for large-scale screenings.
- Ease of sampling
Because one-time screening is not repeatedly done, ease of sampling is a less important criterion.

Result interpretation

Rapid antigen tests have shown to have a lower sensitivity in asymptomatic people and a negative result does therefore not exclude an infection. Rapid antigen tests have a good sensitivity in people with a high viral load and will detect the most infectious infections. Nevertheless, we cannot exclude that some of the missed infections will be infectious. A negative result does therefore not exempt the tested person from respecting all needed preventive and protective measures.

Rapid antigen tests have overall a high specificity. Nevertheless, in settings with a (very) low positivity rate a very high specificity is necessary to avoid too many false positive results. We therefore only recommend one-time screening using a rapid antigen tests with a very high specificity (>99%) and in populations where the expected positivity rate will be at least 1%.

3.5. Recommendations

- One-time screening of visitors to nursing homes is considered a useful strategy, and can optionally be implemented.
- Systematic one-time screening in other situations is only justifiable if the following conditions are fulfilled:
 - Risk of rapid spread of the infection to a large number of people; AND
 - No possibility to fully apply effective protective measures; AND
 - The number of testing events is feasible.
- In situations where an immediate test result is needed (less than 30 minutes), we recommend the use of an Ag RDT with high specificity (>99%) on a naso-pharyngeal or a nasal/oral swab. A PCR test can be used in settings where their result can rapidly be obtained, such as in hospitals.
- If a rapid antigen test is used, a negative result does not exclude infection or infectiousness, and all preventive and protective measures need to be respected.

References

1. Universal screening for SARS-CoV-2 infection: a rapid review - Viswanathan, M - 2020 | Cochrane Library [Internet]. [cited 2020 Dec 10]. Available from: <https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD013718/full>
2. Cochrane Rapid Review investigates the effectiveness of screening for COVID-19 [Internet]. [cited 2020 Dec 10]. Available from: </news/cochrane-rapid-review-investigates-effectiveness-screening-covid-19>

4. USE OF MULTIPLEX PCR ASSAYS

4.1. Current recommendation

The testing strategy update of the RAG on 09/09/20 stated that during the flu season symptomatic patients, especially in severe condition, could be tested with a multiplex PCR panel in order to guide therapeutic care and support:

- Hospitalized patients with respiratory symptoms: multiplex PCR mini panel (SARS-CoV2, RSV, Influenza A and B) on nasopharyngeal or nose/throat sample
- Hospitalized patients with respiratory symptoms and severe condition: multiplex PCR maxi panel (detection of 10 pathogens)
- Symptomatic patients in ambulatory care: only test for SARS-CoV-2

Limitations to the routine use of multiplex PCR highlighted on 09/09/2020 included:

- No commercially available multiplex test in Belgium
- No reimbursement of multiplex test
- Not all labs can perform the multiplex test
- Lab personnel has often no information on the clinical profile (symptomatic or asymptomatic; severity) of the patient

4.2. literature

Multiplex PCRs have been used to analyze transmission patterns of different respiratory pathogens as well as to assess the extent of co-infections of SARS-CoV-2 and other common respiratory pathogens, and its impact on clinical outcomes.

In a recent study, Poole et al. assessed the impact of SARS-CoV-2 in the prevalence of respiratory viruses in hospitalized patients (1). They compared samples from March-May 2020 with samples from the same period in previous years. Before 2020, non-SARS-CoV-2 viruses (such as Influenza, rhinovirus, RSV, seasonal coronaviruses or parainfluenza virus) were detected in 54 % of patients. In 2020 non-SARS-CoV-2 viruses were present in 4,1 % of the samples while 38 % of samples were positive for SARS-CoV-2. The emergence of SARS-CoV-2 was therefore associated with reductions in the circulation of seasonal respiratory viruses. The authors concluded that this observation could be due to the measures taken to fight COVID-19, such as social distancing and lock-down. Another hypothesis points at interactions and interferences between different viruses. This has been shown for other respiratory viruses (2).

Reduction in the circulation of other seasonal respiratory viruses during the first peak of the epidemic was also observed in several regions worldwide (3,4). An early Italian study however did not see different trends for other respiratory viruses in March 2020 compared to the same period in previous years (5).

Co-infections of SARS-CoV-2 and other respiratory viruses have been described in several reports, with variable levels of co-infections. In a study by Leuzinger et al. co-infection was shown to occur in 1,8 % of the samples (6). Poole et al. found co-infections of SARS-CoV-2 and other respiratory virus in 1 % of the samples (1). Also other studies found low levels of coinfection with other respiratory viruses (7). Some studies observed more extended cases of co-infections with bacterial pathogens (8).

COVID-19 patients co-infected with influenza had in one study a 2.27 times greater risk of death than non-co-infected patients (9).

References

1. Poole S, Brendish NJ, Clark TW. SARS-CoV-2 has displaced other seasonal respiratory viruses: Results from a prospective cohort study. *J Infect*. 2020 Nov;S0163445320307076.
2. Nickbakhsh S, Mair C, Matthews L, Reeve R, Johnson PCD, Thorburn F. Virus–virus interactions impact the population dynamics of influenza and the common cold. :9.
3. Marriott D, Beresford R, Mirdad F, Stark D, Glanville A, Chapman S, et al. Concomitant marked decline in prevalence of SARS-CoV-2 and other respiratory viruses among symptomatic patients following public health interventions in Australia: data from St Vincent’s Hospital and associated screening clinics, Sydney, NSW. *Clin Infect Dis Off Publ Infect Dis Soc Am* [Internet]. 2020 Aug 25 [cited 2020 Dec 3]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7499558/>
4. Zhang K, Misra A, Kim PJ, Moghadas SM, Langley JM, Smieja M. Rapid disappearance of influenza following the implementation of COVID-19 mitigation measures in Hamilton, Ontario. *medRxiv*. 2020 Nov 30;2020.11.27.20240036.
5. Sberna G, Amendola A, Valli MB, Carletti F, Capobianchi MR, Bordi L, et al. Trend of respiratory pathogens during the COVID-19 epidemic. *J Clin Virol*. 2020 Aug;129:104470.
6. Leuzinger K, Roloff T, Gosert R, Sogaard K, Naegele K, Rentsch K, et al. Epidemiology of Severe Acute Respiratory Syndrome Coronavirus 2 Emergence Amidst Community-Acquired Respiratory Viruses. *J Infect Dis* [Internet]. 2020 Jul 29 [cited 2020 Dec 3]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7454752/>
7. Wee LE, Ko KKK, Ho WQ, Kwek GTC, Tan TT, Wijaya L. Community-acquired viral respiratory infections amongst hospitalized inpatients during a COVID-19 outbreak in Singapore: co-infection and clinical outcomes. *J Clin Virol Off Publ Pan Am Soc Clin Virol*. 2020;128:104436.
8. Lai C-C, Wang C-Y, Hsueh P-R. Co-infections among patients with COVID-19: The need for combination therapy with non-anti-SARS-CoV-2 agents? *J Microbiol Immunol Infect*. 2020 Aug;53(4):505–12.
9. Stowe J, Tessier E, Zhao H, Guy R, Muller-Pebody B, Zambon M, et al. Interactions between SARS-CoV-2 and Influenza and the impact of coinfection on disease severity: A test negative design. *medRxiv*. 2020 Sep 22;2020.09.18.20189647.

4.3. International recommendations


WHO

The World Health Organization recommends in a Policy brief dated 6 November to test patients with severe or complicated disease or those with risk factors (regardless of severity) using a rapid molecular assay when results can be made available within 24 hours preferably¹⁴. Awaiting test results should not delay empiric treatment, which can be modified subsequently, according to test results. The longer the time lag between sampling and test results, the less the test will benefit clinical management. Empiric treatment without laboratory diagnosis could lead to expanded use of oseltamivir and could contribute to overuse and the development of resistance.

¹⁴ WHO - Readiness for influenza during the COVID-19 pandemic - Policy brief - 6 November 2020. Available at: file:///C:/Users/YvLa1869/Downloads/WHO-2019-nCoV-Influenza_readiness_COVID-19-2020.1-eng.pdf

4.4. Influenza season in Belgium

The figure below presents the trends in influenza like illnesses (ILI) in past years in Belgium. The period with the highest incidence of ILI in general practitioners consultations is from January to April.



4.5. Discussion

- Many of the factors limiting the use of multiplex assays have been resolved. They have become available on the Belgian market, most labs can perform them and lab staff can know what type of multiplex to use from the prescriptions.
- The only persisting barrier is that currently multiplex assays are still not reimbursed by the national health insurance.
- Despite the evidence showing that the emergence of SARS-CoV-2 appears to result in reductions in the circulation of seasonal respiratory viruses and the low levels of co-infection established in several studies, it was agreed that it remains important to use multiplex assays in certain situations. In particular because a co-infection with, for example, influenza can give a more severe pathology. This applies both to adults and children.
- Because of the high cost of a multiplex assay, the use should be reserved for the more severe cases in hospital settings. A broader use in primary care settings is not indicated. Eligible patients include both those requiring hospitalization and those presenting at the emergency department.
- Maxi panels, testing for a broad range of panels, should be reserved for patients with a severe clinical condition. Patients without a severe clinical condition can be tested with a mini panel.
- There are mini panels that only include SARS-CoV-2 and influenza (A and B), and mini panels that also include RSV. Both are acceptable.
- A sequential approach (first test for SARS-CoV-2 and only use a multiplex assay if negative) is not recommended.

- It is not known what the effect of the measures to control the SARS-CoV-2 epidemic will have on the seasonal influenza epidemic. It is therefore important to monitor the incidence of influenza and only initiate the use of multiplex assays once there are clear signs of a seasonal epidemic.

4.6. Updated recommendation for use of multiplex PCR in Belgium

- During the seasonal influenza epidemic patients with respiratory symptoms, especially in severe condition, can be tested with a multiplex PCR panel in order to guide therapeutic care and support.
- The use of multiplex assays is only recommended if there is evidence of a seasonal influenza epidemic, based on the surveillance data of influenza-like illnesses in Belgium or neighboring countries.
- A multiplex assay is only indicated in patients attending hospitals, both hospitalized in-patients and out-patients at the emergency department. Patients in ambulatory care outside a hospital setting (such as in general practice) should only be tested for SARS-CoV-2.
- Patients with respiratory symptoms without severe clinical condition should be tested with a multiplex PCR mini panel (SARS-CoV2/ Influenza A and B, or SARS-CoV2/ Influenza A and B/ RSV).
- Patients with respiratory symptoms and a severe clinical condition can be tested with a multiplex PCR maxi panel (detection of a broad range of pathogens), if available.
- Multiplex assays should be done on specimens that ensure a high sensitivity, such as nasopharyngeal swabs, nose/throat swabs, tracheal aspirate or broncheo-alveolar lavage fluid (BAL); and not on specimens with lower sensitivity, such as salivary samples.